

*Roundtrip Spanners and Roundtrip Routing in
Directed Graphs*

by Roditty, Thorup, and Zwick (2008)

Yuchong Pan

MIT 6.890

December 9, 2021

Spanners in **Undirected** Graphs

Definition (Spanner)

Let $G = (V, E)$ be a weighted **undirected** graph. A subgraph H of G is a t -**spanner** if $d_H(u, v) \leq t \cdot d_G(u, v)$ for all $u, v \in V$.

Spanners in **Undirected** Graphs

Definition (Spanner)

Let $G = (V, E)$ be a weighted **undirected** graph. A subgraph H of G is a t -**spanner** if $d_H(u, v) \leq t \cdot d_G(u, v)$ for all $u, v \in V$.

Theorem (Althöfer et al., 1993)

*For any $k \in \mathbb{N}$, any weighted **undirected** graph on n vertices has a $(2k - 1)$ -spanner with $O(n^{1+1/k})$ edges.*

Spanners in **Undirected** Graphs

Definition (Spanner)

Let $G = (V, E)$ be a weighted **undirected** graph. A subgraph H of G is a t -**spanner** if $d_H(u, v) \leq t \cdot d_G(u, v)$ for all $u, v \in V$.

Theorem (Althöfer et al., 1993)

*For any $k \in \mathbb{N}$, any weighted **undirected** graph on n vertices has a $(2k - 1)$ -spanner with $O(n^{1+1/k})$ edges.*

Directed graphs?

Spanners in **Undirected** Graphs


Definition (Spanner)

Let $G = (V, E)$ be a weighted **undirected** graph. A subgraph H of G is a t -**spanner** if $d_H(u, v) \leq t \cdot d_G(u, v)$ for all $u, v \in V$.

Theorem (Althöfer et al., 1993)

For any $k \in \mathbb{N}$, any weighted **undirected** graph on n vertices has a $(2k - 1)$ -spanner with $O(n^{1+1/k})$ edges.

Directed graphs?



Spanners in **Undirected** Graphs


Definition (Spanner)

Let $G = (V, E)$ be a weighted **undirected** graph. A subgraph H of G is a t -**spanner** if $d_H(u, v) \leq t \cdot d_G(u, v)$ for all $u, v \in V$.

Theorem (Althöfer et al., 1993)

For any $k \in \mathbb{N}$, any weighted **undirected** graph on n vertices has a $(2k - 1)$ -spanner with $O(n^{1+1/k})$ edges.

Directed graphs?



Roundtrip Spanners

Definition (Roundtrip spanner)

Let G be a weighted **directed** graph. A subgraph H of G is a **t -roundtrip-spanner** if $d_H(u \rightleftarrows v) \leq t \cdot d_G(u \rightleftarrows v)$ for $u, v \in V$.

Roundtrip Spanners

Definition (Roundtrip spanner)

Let G be a weighted **directed** graph. A subgraph H of G is a **t -roundtrip-spanner** if $d_H(u \rightleftarrows v) \leq t \cdot d_G(u \rightleftarrows v)$ for $u, v \in V$.

Theorem (Cowen and Wagner, 2000)

*For any $k \in \mathbb{N}$, any weighted **directed** graph on n vertices has a $(2^k - 1)$ -roundtrip-spanner with $\tilde{O}(n^{1+1/k})$ edges.*

Roundtrip Spanners

Definition (Roundtrip spanner)

Let G be a weighted **directed** graph. A subgraph H of G is a **t -roundtrip-spanner** if $d_H(u \rightleftarrows v) \leq t \cdot d_G(u \rightleftarrows v)$ for $u, v \in V$.

Theorem (Cowen and Wagner, 2000)

For any $k \in \mathbb{N}$, any weighted **directed** graph on n vertices has a $(2^k - 1)$ -roundtrip-spanner with $\tilde{O}(n^{1+1/k})$ edges.

Theorem (Roditty, Thorup, and Zwick, 2008)

For any $k \in \mathbb{N}, \varepsilon > 0$, any weighted **directed** graph on n vertices with edge weights from $[1, W]$ has a $(2k + \varepsilon)$ -roundtrip-spanner with $O((k^2/\varepsilon)n^{1+1/k} \log(nW))$ edges.

A General Technique of Cohen (1998)

A General Technique of Cohen (1998)

Procedure PARTIALSPANNER(R)

Input: $R \in [1, 2nW]$.

Output: A subgraph H of G with $O(kn^{1+1/k})$ edges such that $d_H(u \rightleftharpoons v) \leq 2kR$ for all $u, v \in V$ with $d_G(u \rightleftharpoons v) \leq R$.

A General Technique of Cohen (1998)

Procedure PARTIALSPANNER(R)

Input: $R \in [1, 2nW]$.

Output: A subgraph H of G with $O(kn^{1+1/k})$ edges such that $d_H(u \rightleftharpoons v) \leq 2kR$ for all $u, v \in V$ with $d_G(u \rightleftharpoons v) \leq R$.

The Approach of Cohen (1998)

```
1 foreach  $i \leftarrow 1, \dots, \log_{1+\epsilon}(2nW)$  do  
2    $R_i \leftarrow (1 + \epsilon)^i$   
3    $H_i \leftarrow \text{PARTIALSPANNER}(R_i)$   
4  $H \leftarrow \bigcup \{H_i : i \in [\log_{1+\epsilon}(2nW)]\}$   
5 return  $H$ 
```

A General Technique of Cohen (1998)

Procedure PARTIALSPANNER(R)

Input: $R \in [1, 2nW]$.

Output: A subgraph H of G with $O(kn^{1+1/k})$ edges such that $d_H(u \rightleftharpoons v) \leq 2kR$ for all $u, v \in V$ with $d_G(u \rightleftharpoons v) \leq R$.

The Approach of Cohen (1998)

```
1 foreach  $i \leftarrow 1, \dots, \log_{1+\varepsilon}(2nW)$  do  
2    $R_i \leftarrow (1 + \varepsilon)^i$   
3    $H_i \leftarrow \text{PARTIALSPANNER}(R_i)$   
4  $H \leftarrow \bigcup \{H_i : i \in [\log_{1+\varepsilon}(2nW)]\}$   
5 return  $H$ 
```

Claim: H is a $2k(1 + \varepsilon)$ -roundtrip-spanner of G with $O(kn^{1+1/k} \log_{1+\varepsilon}(nW))$ edges.

Constructing Partial Spanners

Definition (Ball)

For every $V' \subset V$, $v \in V$ and $r \geq 0$, let

$$\text{ball}_{V'}(v, r) = \{u \in V' : d_{G[V']}(v \rightleftharpoons u) \leq r\},$$

called a **ball** centered at v of radius r .

Constructing Partial Spanners

Definition (Ball)

For every $V' \subset V$, $v \in V$ and $r \geq 0$, let

$$\text{ball}_{V'}(v, r) = \{u \in V' : d_{G[V']}(v \rightleftarrows u) \leq r\},$$

called a **ball** centered at v of radius r .

Definition ((k, R) -cover)

A set \mathcal{C} of balls is a (k, R) -**cover** of a directed graph $G = (V, E)$ if each ball in \mathcal{C} is of radius at most kR , and for all $u, v \in V$ with $d_G(u \rightleftarrows v) \leq R$, there is a ball $B \in \mathcal{C}$ such that $u, v \in B$.

Constructing Partial Spanners

Procedure COVER(G, k, R)

```
1  $\mathcal{C} \leftarrow \emptyset$ 
2  $V_{k-1} \leftarrow V(G)$ 
3 foreach  $i \leftarrow k-1, \dots, 0$  do
4    $S_i \leftarrow \text{SAMPLE}(V_i, n^{-i/k})$ 
5    $\mathcal{C} \leftarrow \mathcal{C} \cup \{\text{ball}_{V_i}(v, (i+1)R) : v \in S_i\}$ 
6    $V_{i-1} \leftarrow V_i \setminus \bigcup_{v \in S_i} \text{ball}_{V_i}(v, iR)$ 
7 return  $\mathcal{C}$ 
```


Constructing Partial Spanners

Procedure COVER(G, k, R)

```
1  $\mathcal{C} \leftarrow \emptyset$ 
2  $V_{k-1} \leftarrow V(G)$ 
3 foreach  $i \leftarrow k-1, \dots, 0$  do
4    $S_i \leftarrow \text{SAMPLE}(V_i, n^{-i/k})$ 
5    $\mathcal{C} \leftarrow \mathcal{C} \cup \{\text{ball}_{V_i}(v, (i+1)R) : v \in S_i\}$ 
6    $V_{i-1} \leftarrow V_i \setminus \bigcup_{v \in S_i} \text{ball}_{V_i}(v, iR)$ 
7 return  $\mathcal{C}$ 
```

Theorem

*The collection $\mathcal{C} = \text{COVER}(G, k, R)$ is a (k, R) -cover of a directed graph $G = (V, E)$. For each $v \in V$, the **expected** number of balls in \mathcal{C} containing v is at most $kn^{1/k}$.*

COVER(G, k, R) Is a (k, R) -Cover

- Let $u, v \in V$ be such that $d_G(u \rightleftarrows v) \leq R$.
- Let c be a closed tour in G of length $\leq R$ containing u and v .
- Let i be the largest index so that a vertex u' on c is contained in the **core** $\text{ball}_{V_i}(w, iR)$ of $\text{ball}_{V_i}(w, (i+1)R)$ added to \mathcal{C} .
- By maximality of i , all the vertices on c are contained in V_i .
- By the triangle inequality,

$$\begin{aligned}d_{G[V_i]}(w \rightleftarrows v) &\leq d_{G[V_i]}(w \rightleftarrows u) + d_{G[V_i]}(u \rightleftarrows v) \\ &\leq iR + R = (i+1)R.\end{aligned}$$

- Therefore, $v \in \text{ball}_{V_i}(w, (i+1)R)$.
- Similarly, $u \in \text{ball}_{V_i}(w, (i+1)R)$, completing the proof.

Expected Number of Balls Containing a Vertex

Lemma

In each iteration, the expected number of balls containing $v \in V$ that are added to \mathcal{C} is at most $n^{1/k}$.


Expected Number of Balls Containing a Vertex

Lemma

In each iteration, the expected number of balls containing $v \in V$ that are added to \mathcal{C} is at most $n^{1/k}$.

- Let $B = \text{ball}_{V_{i+1}}(v, (i+1)R)$.
- If $u \in B$ is chosen to S_{i+1} (with probability $n^{-(i+1)/k}$), then $v \in \text{ball}_{V_{i+1}}(u, (i+1)R)$ and thus removed in iteration $i+1$.
- If $v \in \text{ball}_{V_i}(u, (i+1)R) \subset \text{ball}_{V_{i+1}}(u, (i+1)R)$, then $u \in B$.
- $\mathbb{P}[u \in S_i \wedge v \in \text{ball}_{V_i}(u, (i+1)R)] = (1 - n^{-(i+1)/k})^{|B|} n^{-i/k}$.
- The expected number of balls containing v in iteration i is $|B| (1 - n^{-(i+1)/k})^{|B|} n^{-i/k}$.

Expected Number of Balls Containing a Vertex



$\mathbb{E}[\# \text{ balls containing } v \text{ in iteration } i]$

$$= |B| \left(1 - n^{-(i+1)/k}\right)^{|B|} n^{-i/k} \leq \frac{n^{-i/k}}{en^{-(i+1)/k}} = \frac{n^{1/k}}{e}.$$

Double-Trees

Definition (Double-tree)

Let $G = (V, E)$ be a weighted directed graph. Let $V' \subset V$, $v \in V$ and $r \geq 0$. Let $B = \text{ball}_{V'}(v, r)$.

Let $\text{OutTree}(B, v)$ be a tree containing directed shortest paths in $G[V']$ from v to all the vertices in B .

Let $\text{InTree}(B, v)$ be a tree containing directed shortest paths in $G[V']$ from all the vertices in B to v .

Let $\text{InOutTrees}(B, v) = \text{InTree}(B, v) \cup \text{OutTree}(B, v)$, referred to as a **double-tree**.

Double-Trees

Definition (Double-tree)

Let $G = (V, E)$ be a weighted directed graph. Let $V' \subset V$, $v \in V$ and $r \geq 0$. Let $B = \text{ball}_{V'}(v, r)$.

Let $\text{OutTree}(B, v)$ be a tree containing directed shortest paths in $G[V']$ from v to all the vertices in B .

Let $\text{InTree}(B, v)$ be a tree containing directed shortest paths in $G[V']$ from all the vertices in B to v .

Let $\text{InOutTrees}(B, v) = \text{InTree}(B, v) \cup \text{OutTree}(B, v)$, referred to as a **double-tree**.

Lemma

Let $V' \subset V$ and $v \in V$. If $u \in B := \text{ball}_{V'}(v, r)$ and w is on a shortest path in $G[V']$ from u to v , or from v to u , then $w \in B$.

Double-Trees

Definition (Double-tree)

Let $G = (V, E)$ be a weighted directed graph. Let $V' \subset V$, $v \in V$ and $r \geq 0$. Let $B = \text{ball}_{V'}(v, r)$.

Let $\text{OutTree}(B, v)$ be a tree containing directed shortest paths in $G[V']$ from v to all the vertices in B .

Let $\text{InTree}(B, v)$ be a tree containing directed shortest paths in $G[V']$ from all the vertices in B to v .

Let $\text{InOutTrees}(B, v) = \text{InTree}(B, v) \cup \text{OutTree}(B, v)$, referred to as a **double-tree**.

Lemma

Let $V' \subset V$ and $v \in V$. If $u \in B := \text{ball}_{V'}(v, r)$ and w is on a shortest path in $G[V']$ from u to v , or from v to u , then $w \in B$.

Corollary

$V(\text{InOutTrees}(B, v)) \subset B$, so $e(\text{InOutTrees}(B, v)) \leq 2(|B| - 1)$.

Double-Trees

Lemma

Let $V' \subset V, v \in V$. If $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$, $\text{InOutTrees}(B, v)$ contains a closed directed tour containing u_1, u_2 of length $\leq 2r$.

Double-Trees

Lemma

Let $V' \subset V, v \in V$. If $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$, $\text{InOutTrees}(B, v)$ contains a closed directed tour containing u_1, u_2 of length $\leq 2r$.

Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of G :

Double-Trees

Lemma

Let $V' \subset V, v \in V$. If $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$, $\text{InOutTrees}(B, v)$ contains a closed directed tour containing u_1, u_2 of length $\leq 2r$.

Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of G :

- Let $\varepsilon' = \varepsilon/(2k)$.
- For $i \in [\log_{1+\varepsilon'}(2nW)]$, let $\mathcal{C}_i = \text{COVER}(G, k, R_i)$ be a (k, R_i) -cover, where $R_i = (1 + \varepsilon')^i$.
- Let $H = \bigcup \{\text{InOutTrees}(B) : B \in \bigcup_i \mathcal{C}_i\}$.
- Therefore, H is a $(2k + \varepsilon)$ -roundtrip-spanner of G with $O((k^2/\varepsilon)n^{1+1/k} \log(nW))$ edges **in expectation**.

Double-Trees

Lemma

Let $V' \subset V, v \in V$. If $u_1, u_2 \in B := \text{ball}_{V'}(v, r)$, $\text{InOutTrees}(B, v)$ contains a closed directed tour containing u_1, u_2 of length $\leq 2r$.

Obtaining a $(2k + \varepsilon)$ -roundtrip-spanner of G :

- Let $\varepsilon' = \varepsilon/(2k)$.
- For $i \in [\log_{1+\varepsilon'}(2nW)]$, let $\mathcal{C}_i = \text{COVER}(G, k, R_i)$ be a (k, R_i) -cover, where $R_i = (1 + \varepsilon')^i$.
- Let $H = \bigcup \{\text{InOutTrees}(B) : B \in \bigcup_i \mathcal{C}_i\}$.
- Therefore, H is a $(2k + \varepsilon)$ -roundtrip-spanner of G with $O((k^2/\varepsilon)n^{1+1/k} \log(nW))$ edges **in expectation**.

Application: Compact roundtrip routing schemes.

State-of-the-Art

Theorem (Cen, Duan, and Gu, 2019)

For any $k \in \mathbb{N}$, any weighted directed graph on n vertices has a $(2k - 1)$ -roundtrip-spanner with $O(kn^{1+1/k} \log n)$ edges.

State-of-the-Art

Theorem (Cen, Duan, and Gu, 2019)

For any $k \in \mathbb{N}$, any weighted directed graph on n vertices has a $(2k - 1)$ -roundtrip-spanner with $O(kn^{1+1/k} \log n)$ edges.

Proposition

*If the Erdős girth conjecture is true, there is an **undirected** graph on n vertices such that any $(2k - 1)$ -spanner has $\Omega(kn^{1+1/k})$ edges.*

Thank you and happy holidays!